




Individual-days




Aggressive acts to self




Aggressive acts to others




Individuals having alleged abuse, neglect or exploitation




Body mass index




Individuals with body mass index greater than 25 (overweight to obese)




Body mass index change




Weight change




Waist circumference, males




Combined pharmacotherapy




Electroconvulsive treatment (Each reported as zero for each of the months reported)




Escape/Walkaway




Falls




Homicide




**Incidents of individuals testing positive for
illicit substance use**
(Reported as zero incidents for each month)




Medication variance (estimated)




Medication variance (estimated)




Medication variance (estimated)




Medication variance (estimated)




Any event involving a medication error which results in a major injury or exacerbation of a disease or disorder




Mortality




Non-adherence to WRP for more than 20% of the interventions in seven consecutive days



Close observation




Aspiration pneumonia




Bowel Dysfunction


Individuals with occurrences of fecal impaction or bowel obstruction




Individuals with diagnosis of diabetes mellitus (estimated)




Individuals with diagnosis of diabetes mellitus (estimated)




Individuals with diagnosis of diabetes mellitus (estimated)



Dysphagia




Fractures



Individuals with diagnosis of hepatitis C (estimated)




Hospitalizations




Individuals with diagnosis of MRSA


(Note: As of September 2007, count includes individuals whose infections have healed but who have carrier status.)




Individuals with diagnosis of osteoporosis (estimated)




Individuals with diagnosis of polydipsia (estimated)




Seizure disorder




Individuals receiving older anticonvulsants




Skin integrity




PRN medications




Restraint



Seclusion



Stat medications



Suicide

